


## EE 3025 S2005 Homework Set #3

(due 10:10 AM Friday, February 11, 2005)

**Directions:** Work all 5 problems. We will grade Problem 1 and will randomly choose two of the other problems for grading.

1. Consider the relay circuit in the following diagram:



Relay switches  $A, B, C, D, E, F, G, H$  operate independently; each of them has the same probability  $p$  of working correctly. Let  $P(1 \rightarrow 2)$  denote the probability that a connection will be made from point 1 to point 2. Obviously,  $P(1 \rightarrow 2)$  will be a complicated function of  $p$ . In this problem, you are not going to find the explicit formula for  $P(1 \rightarrow 2)$  as a function of  $p$ . Instead, you will be using Matlab simulations of the circuit to obtain estimates of  $P(1 \rightarrow 2)$  for various  $p$  values. Before starting on this Matlab problem, you might want to study Experiment 3 of Recitation 2.

- (a) Use Matlab to do 50000 simulated trials of the circuit with  $p = 0.90$ . Estimate  $P(1 \rightarrow 2)$  based on these trials. You should run your 50000 trials more than once to see if your estimate is “robust” (that is, hopefully your estimate will not change very much).
  - (b) By trial and error, do 50000 simulated trials over and over again with different values of  $p$  on each run of 50000, until you find a value of  $p$  for which the estimate for  $P(1 \rightarrow 2)$  is as close to 0.85 as you can get.
2. This problem, as part of its solution, will involve the binomial distribution in some way.
- (a) Flip 10 fair dies. Compute the probability that at least 7 of the dies yield a number  $\geq 5$ .
  - (b) Select 10 light bulbs and let each of them burn until they burn out. Each light bulb has a lifetime which is exponentially distributed with mean lifetime 1000 hours. Compute the probability that between 3 and 6 of the light bulbs (inclusively) burn for at least 1200 hours.

3. Dropout is a kind of defect in magnetic tape in which the signal disappears for a brief period of time. We assume that the number of dropouts in a fixed length of tape is a Poisson random variable.

(a) Suppose the number of dropouts in an 1800ft length of tape is a Poisson RV with parameter 3. What is the probability of 3 or more dropouts in this length of tape? What is the conditional probability of 3 or more dropouts given that there is at least one dropout?

(b) Suppose again that the number of dropouts in an 1800ft length of tape is a Poisson RV with parameter 3. Let random variable  $X$  be the number of dropouts in a 3600ft of tape.  $X$  is a Poisson random variable. What is the parameter? Compute the probability that  $X \geq 6$ .

4. A mixed random variable  $X$  has probability density function

$$f_X(x) = (1/6)\delta(x - 5) + (1/6)\delta(x - 10) + (1/6)\delta(x - 15) + \frac{1}{20\sqrt{2\pi}} \exp\left(-\frac{(x - 20)^2}{200}\right),$$

defined for all real  $x$ .

(a) Attach a Matlab plot of the cumulative distribution function  $F_X(x)$  over the range  $0 \leq x \leq 40$ .

(b) Compute  $P(7.5 \leq X \leq 30)$ .

(c) Given that  $X$  is between 7.5 and 30, compute the conditional probability that  $X \geq 12.5$ .

5. This problem regards the computations of means and variances of random variables.

(a) A discrete RV  $X$  takes the values 1, 2, 3, 4, 5 and its PMF takes the form

$$p_X(x) = Cx^2, \quad x = 1, 2, 3, 4, 5,$$

where  $C$  is a positive constant. Compute  $C$ , and then compute the mean and variance of  $X$ .

(b) Now we take  $X$  to be a continuous RV with PDF of the form

$$f_X(x) = Cx^2, \quad 1 \leq x \leq 5 \text{ (zero elsewhere)},$$

where  $C$  is a positive constant. Compute  $C$ , and then compute the mean and variance of  $X$ .

**Supplementary Problems:** (not to hand in) From the textbook, you can try Problems 2.3.2, 2.3.4, 2.4.1, 3.3.6(a)(b), 3.5.1